

# New taxa and new combinations in Western Australian Pittosporaceae

By Eleanor M. Bennett

## Abstract

It is proposed that the genera *Marianthus* and *Billardiera* be combined under *Billardiera*. The necessary new combinations are made, and three new varieties described. Two new varieties are described in the genus *Cheiranthera*.

A new combination, using the earliest available epithet, is provided for the solitary species of *Pronaya*.

## Introduction

The genera *Marianthus* and *Billardiera* are here combined under *Billardiera*. These genera have been distinguished by former workers on their fruit which in *Billardiera* is a berry and in *Marianthus* is a capsule. This division does not appear to be a natural one since a separation based on floral characters would provide a quite different arrangement of species. The use of fruit characters for delimiting these two genera also makes impossible the ready identification of flowering material, as an examination of the ovary provides no clue to its future form and in fact several species have been, and continue to be, incorrectly placed.

In addition two new varieties of *Cheiranthera* are described, one in each of the two species which occur in Western Australia.

In *Pronaya*, the name *P. elegans* Hueg., which has been in common use, is antedated by the name *Spiranthera fraseri* Hook. and a new combination is made accordingly.

Species have been accepted after consulting type specimens, descriptions and dates of publication, occasionally resulting in the changing of previously accepted names.

These new combinations and varieties are being published prior to a revision of the Western Australian species of *Pittosporaceae* in a forthcoming Flora part.

## Billardiera

**Billardiera** Sm., Spec. Bot. Nov. Holl. 1:t.1 (1793). *Type*: *B. scandens* Sm.

*Labillardiera* Roem. et Schult., Syst. Veg. 5:330 (1819). nom. illeg. based on above.

*Marianthus* Hueg. ex Endl., Enum. Pl. Hueg. 8 (1837). *Type*: *M. candidus* Hueg. ex Endl.

*Oncosporum* Putterl. in Endl. et Fenzl., Nov. Stirp. Dec. 9 (1839). *Type*: *C. bicolor* Putterl.

*Calopetalon* J. Drumm. ex Harv., Hooker's Journ. Bot. Kew Gard. Misc. 7:52 (1855). *Type*: *C. ringens* J. Drumm. ex Harv.

*Cyathomiscus* Turcz., Bull. Soc. Imp. Nat. Mosc. 36/1:562 (1863). *Type*: *C. cuneatum* Turcz.

**Billardiera laxiflora** (Benth.) E. M. Bennett comb. nov.

*Marianthus* ? *laxiflorus* Benth., Fl. Austral. 1:119 (1863). *Lectotype*: Herb. Mueller, Porongurup Range, W. Australia, (holo:MEL).

*Billardiera gracilis* Diels, Bot. Jahrb. 35:213 (1904). *Type*: King George Sound, Diels 2259 and Pritzel 252 (syn:PERTH).

**Billardiera villosa** (Turcz.) E. M. Bennett comb. nov.

*Oncosporum villosum* Turcz., Bull. Soc. Imp. Nat. Mosc. 27:365 (1854).

Type: Western Australia, *Drummond* 5th Coll., n.242 (iso:MEL).

*Oncosporum microphyllum* Turcz., loc. cit. Type: not given.

*Marianthus microphyllum* (Turcz.) Benth., Fl. Austral. 1:117 (1863).

*M. rhytidosporus* F. Muell., Fragm. 2:145 (1861). Type: Near Esperance Bay, W. Australia, *Maxwell* (holo:MEL).

**Billardiera granulata** (Turcz.) E. M. Bennett comb. nov.

*Oncosporum granulatum* Turcz., Bull. Soc. Imp. Nat. Mosc. 27:366 (1854).

Type: Western Australia, *Drummond* 3rd Coll., n.210 (iso:W).

*Marianthus granulatus* (Turcz.) Benth., Fl. Austral. 1:118 (1863).

*M. parviflorus* F. Muell., Fragm. 2:144 (1861). Type: Porongurup Range, W. Australia, *Maxwell* (holo:MEL).

**Billardiera candida** (Hueg. ex Endl.) E. M. Bennett comb. nov.

*Marianthus candidus* Hueg. ex Endl., Enum. Pl. Hueg. 8 (1837).

Type: Western Australia, Huegel (holo:W).

**Billardiera coeruleo-punctata** (Klotsch) E. M. Bennett comb. nov.

*Marianthus coeruleo-punctatus* Klotsch in Link, Klotsch et Otto, Ic. Pl. Rar. 28 t. 12 (1840). Lectotype: Western Australia, 1839, *Drummond* (iso:W).

*Pronaya pedunculosa* Turcz., Bull. Soc. Nat. Mosc. 36:560 (1863).

Type: Western Australia, 1839, J. *Drummond* (holo:W).

**Billardiera drummondiana** (Putterl.) E. M. Bennett comb. nov.

*Oncosporum drummondianum* Putterl. in Lehm., Pl. Preiss. 1:194 (1844).

Lectotype: Western Australia, Preiss 1288 (holo:W).

*Marianthus drummondianus* (Putterl.) Benth., Fl. Austral. 1:119 (1863).

*Billardiera dorrienii* Domin. Mém. Soc. Sc. Bohême 2:24 (1923) (holo:K, photo seen).

**B. drummondiana** var. **collina** E. M. Bennett var. nov. (L. *collinus*—occurring on hills)

Folia, caules, calyces et pedicelli dense hirsuti pilis longis unicellulosis etiam breves multicellulosis. Pedicelli 4–10 (15) mm longi.

Type: Bushmead-Kalamunda area, W. Australia, 19 Sep. 1966, F. W. Humphreys 24, (holo:PERTH).

Leaves stems, calyces and pedicels densely hirsute with long unicellular and short multicellular hairs. Pedicels 4–10(15) mm long.

The typical variety differs from var. *collina* in having only unicellular hairs scattered on the leaves, stems and calyces and in having long pedicels. It is found around Cranbrook and Manjimup while var. *collina* is found in the Darling Ranges.

**Billardiera bicolor** (Putterl.) E. M. Bennett comb. nov.

*Oncosporum bicolor* Putterl. in Endl. et Fenzl., Nov. Stirp. Dec. 9 (July 1839). Type: Western Australia, Huegel (n.v.).

*Marianthus pictus* Lindl., Sketch Veg. Swan Riv. Col. 22 (Dec. 1839).

Type: Western Australia (holo:CGE, photo seen).

**B. bicolor** var. **lineata** (F. Muell.) E. M. Bennett comb. et stat. nov.

*M. lineatus* F. Muell., Fragm. 1:27 (1859). Type: Murchison River, W. Australia, F. Mueller. (holo:MEL; iso:W).

In var. *lineata*, the pedicel is stout, the corymbs are shorter than or just exceed the leaves and the capsule is cartilaginous. The typical variety has a slender pedicel, corymbs which exceed the leaves and a crustaceous capsule.

**Billardiera erubescens** (Putterl.) E. M. Bennett comb. nov.

*Marianthus erubescens* Putterl. in Endl. et Fenzl., Nov. Stirp. Dec. 60 (1839). Type: Western Australia, 1837, Roe (holo:W).

*M. purpureus* Turcz., Bull. Soc. Nat. Mosc. 27:364 (1854). Type: Western Australia, J. Drummond 4th Coll., n.96, (n.v.).

**Billardiera ringens** (Drumm. ex Harv.) E. M. Bennett comb. nov.

*Calopetalon ringens* Drumm. ex Harv., Hooker's Journ. Bot. Kew Garden. Misc. 7:52 (1855). Type: Chapman, W. Australia, J. Drummond (n.v.).  
*Marianthus ringens* (Drumm. ex Harv.) F. Muell., Fragm. 1:218 (1859).

**Billardiera sericea** (Turcz.) E. M. Bennett comb. nov.

*Pronaya sericea* Turcz., Bull. Soc. Imp. Nat. Mosc. 27:363 (1854).  
Type: Western Australia, J. Drummond 4th Coll., n.97 (iso:MEL).

**Billardiera parviflora** DC., Prodr. 1:346 (1824).

Type: Western Australia, *Leschenault* (holo:G-DC).

*Marianthus temuis* Benth., Fl. Austral. 1:119 (1863). Syntypes: Flinders Bay, W. Australia, Collie (K, MEL); Geographe Bay, *Leschenault* (n.v.); Cape Naturaliste, *Oldfield* (n.v.).

*M. gracilis* Ostenf., Dansk. Vidensk. Selsk. Biol. Medd. 312:67 Pl. 9. (1921). Type: Yallingup Cave, 26 Sept. 1914, *Ostenfeld* 1018. (n.v.).

**B. parviflora** var. **guttata** E. M. Bennett var. nov. (*L. guttatus*—spotted)

Petala caesia, sauce atro-cyaneo-maculata.

Type: Armadale-Jarrahdale, W. Australia, 3 Aug. 1960, F. C. J. Lullfitz (holo:PERTH).

The var. *guttata* differs from the typical variety in the petals having dark blue spots in the throat. It occurs in the southern areas of the Darling Range whereas the typical variety is found near Augusta.

### Cheiranthera

**Cheiranthera preissiana** Putterl. in Lehm., Pl. Preiss. 1:201 (1845).

Type: York, *Preiss* 1291 (iso:MEL).

*C. parviflora* Benth., Fl. Austral. 1:128 (1863). Type: Western Australia, J. Drummond n.34 and 80 (syn:MEL).

**C. preissiana** var. **planifolia** E. M. Bennett var. nov. (*L. planus, folium*—flat leaved)

Folia linearia, plana, glabra vel pilis sparsis. Surculi et folia juvenes pilis sparsis.

Type: Nornalup, W. Australia, E. M. Bennett 3184, 19 Jan. 1970 (holo:PERTH).

Leaves linear, flat, glabrous or with scattered hairs. Young shoots and leaves with scattered hairs.

The typical variety differs from var. *planifolia* in having ovate, pubescent leaves.

**Cheiranthera filifolia** Turcz., Bull. Soc. Nat. Mosc. 27:364 (1854).

Type: Western Australia, *Drummond* 4th Coll. n.94, (iso:MEL).

*C. tortilis* F. Muell., Fragm. 2:79 (1860). Type: South-west of Western Australia, *F. Mueller* (n.v.).

- C. filifolia** var. **brevifolia** (F. Muell.) Benth., Fl. Austral. 1:128 (1863).  
*C. brevifolia* F. Muell., Fragm. 1:97 (1858). *Type:* Phillips Range,  
F. Mueller (holo: MEL).
- C. filifolia** var. **simplicifolia** E. M. Bennett var. nov. (L. *simplex*, *folium*—  
simple leaved)  
Folia plana, linearia, acuminata.  
*Type:* Pindar, W. Australia, C. A. Gardner 7766 (holo: PERTH).  
Leaves flat, linear, acuminate.  
This variety differs from the typical variety and from var. *brevifolia* (F. Muell.) Benth. in having flat leaves. The other two varieties have terete, canaliculate leaves.

### Pronaya

**Pronaya fraseri** (Hook.) E. M. Bennett comb. nov.

*Spiranthera fraseri* Hook., Bot. Mag. 63, sub tab. 3253 (1836).

*Type:* Western Australia, Fraser (n.v.).

*Campylanthera fraseri* (Hook.) Hook., Icones Pl. 1: tab. 82 (Mar. 1837).

*Pronaya speciosa* Endl., in Endl. et al., Enum. Pl. Huegel: 9 (April 1837).

*Type:* King George Sound, W. Australia, Ferd. Bauer (holo: W).

*Pronaya elegans* Hueg., Bot. Archiv. tab. 6 (April-May 1837).

*Type:* Western Australia, F. Bauer (n.v.). *Billardiera elegans* (Hueg.) F. Muell., Pl. Indig. Col. Vict. 1: 78 (1862).

**P. fraseri** var. **minor** (Benth.) E. M. Bennett comb. nov.

*Pronaya elegans* Hueg. var. *minor* Benth., Fl. Austral. 1: 126 (1863).

*Type:* South coast (W. Australia), R. Brown (n.v.).